

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## *Shajia*, a new genus of polyconitid rudist from the Langshan Formation of the Lhasa Block, Tibet, and its palaeogeographical implications

### Journal Item

#### How to cite:

Rao, Xin; Skelton, Peter W.; Sano, Shin-ichi; Zhang, Yiqiang; Zhang, Yichun; Pan, Yanhong; Cai, Huawei; Peng, Bo; Zhang, Tong and Ma, Zhaoxiong (2020). *Shajia*, a new genus of polyconitid rudist from the Langshan Formation of the Lhasa Block, Tibet, and its palaeogeographical implications. *Cretaceous Research*, 105, article no. 104151.

For guidance on citations see [FAQs](#).

© 2019 Elsevier Ltd.



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1016/j.cretres.2019.05.009>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.


---

[oro.open.ac.uk](http://oro.open.ac.uk)

# Accepted Manuscript

*Shajia*, a new genus of polyconitid rudist from the Langshan Formation of the Lhasa Block, Tibet, and its palaeogeographical implications

Xin Rao, Peter W. Skelton, Shin-ichi Sano, Yiqiang Zhang, Yichun Zhang, Yanhong Pan, Huawei Cai, Bo Peng, Tong Zhang, Zhaoxiong Ma



PII: S0195-6671(18)30350-1

DOI: <https://doi.org/10.1016/j.cretres.2019.05.009>

Reference: YCRES 4151

To appear in: *Cretaceous Research*


Received Date: 31 August 2018

Revised Date: 18 January 2019


Accepted Date: 22 May 2019

Please cite this article as: Rao, X., Skelton, P.W., Sano, S.-i., Zhang, Y., Zhang, Y., Pan, Y., Cai, H., Peng, B., Zhang, T., Ma, Z., *Shajia*, a new genus of polyconitid rudist from the Langshan Formation of the Lhasa Block, Tibet, and its palaeogeographical implications, *Cretaceous Research*, <https://doi.org/10.1016/j.cretres.2019.05.009>.


This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



★ *Horiopleura haydeni*



★ *Shajia tibetica* gen. et sp. nov.



1 ***Shajia*, a new genus of polyconitid rudist from the Langshan Formation of the**  
2 **Lhasa Block, Tibet, and its palaeogeographical implications**

3 Xin Rao<sup>1,2\*</sup>, Peter W. Skelton<sup>3\*</sup>, Shin-ichi Sano<sup>4</sup>, Yiqiang Zhang<sup>2</sup>, Yichun Zhang<sup>1</sup>,  
4 Yanhong Pan<sup>5</sup>, Huawei Cai<sup>5</sup>, Bo Peng<sup>6</sup>, Tong Zhang<sup>7</sup>, Zhaoxiong Ma<sup>7</sup>

5 <sup>1</sup>State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of  
6 Geology and Palaeontology and the Center for Excellence in Life and  
7 Paleoenvironment, Chinese Academy of Sciences, 39 East Beijing Road, Nanjing  
8 210008, China; e-mail: [xinrao@nigpas.ac.cn](mailto:xinrao@nigpas.ac.cn), [yczhang@nigpas.ac.cn](mailto:yczhang@nigpas.ac.cn)

9 <sup>2</sup>Department of Invertebrate Palaeontology, Nanjing Institute of Geology and  
10 Palaeontology, Chinese Academy of Sciences, 39 East Beijing Road, Nanjing 210008,  
11 China; e-mail: [yiqiangzhang81@gmail.com](mailto:yiqiangzhang81@gmail.com)

12 <sup>3</sup>School of Environment, Earth and Ecosystem Sciences, The Open University, Milton  
13 Keynes MK7 6AA, UK; e-mail: [pwskelto@waitrose.com](mailto:pwskelto@waitrose.com)

14 <sup>4</sup>Department of Earth System Science, Graduate School of Science and Engineering  
15 for Research, University of Toyama 3190 Gofuku, Toyama-shi, Toyama 930-8555,  
16 Japan; e-mail: [ssano@sus.u-toyama.ac.jp](mailto:ssano@sus.u-toyama.ac.jp)

17 <sup>5</sup>CAS Key Laboratory of Economic Stratigraphy and Palaeogeography, Nanjing  
18 Institute of Geology and Palaeontology and Center for Excellence in Life and  
19 Paleoenvironment, Chinese Academy of Sciences, Nanjing 210008, China; e-mail:  
20 [yhpan@nigpas.ac.cn](mailto:yhpan@nigpas.ac.cn), [hwcai@nigpas.ac.cn](mailto:hwcai@nigpas.ac.cn)

21 <sup>6</sup>Nanjing Center, China Geological Survey, Nanjing 210016, China; e-mail:

22 [pengbocc@163.com](mailto:pengbocc@163.com)

23 <sup>6</sup>Sichuan Geological Survey, No. 25 North Renmin Road, Chengdu 610084, China;

24 e-mail: [986573254@qq.com](mailto:986573254@qq.com), [mzx0120@163.com](mailto:mzx0120@163.com)

25 \*Corresponding authors

## 26 **Abstract.**

27 A new polyconitid rudist *Shajia tibetica* gen. et sp. nov., of late Aptian to Albian  
28 age, is described from the Langshan Formation of Nyima County, northern Lhasa  
29 Block, Tibet. Though comparable in size and external morphology with *Horiopleura*  
30 *haydeni* Douvillé, which is a common endemic species in southwestern Asia, *Shajia*  
31 differs from the latter species in its possession of an inwardly inclined, instead of  
32 outwardly facing, posterior myophore in the right valve. In addition, a single  
33 specimen from Ladakh, which was previously assigned to *Polyconites?* sp., on  
34 account of a similar myophoral distinction from *H. haydeni*, is transferred to the new  
35 genus. *Shajia* is considered most likely to have been derived from one of a group of  
36 *Horiopleura* species that lived on the southern margin of the Mediterranean Tethys.

37 The so-called ‘Yasin fauna’ represented by the late Aptian to Albian *Horiopleura*  
38 *haydeni*/*Auroradiolites gilgitensis* rudist association, is considered to be restricted to  
39 southwestern Asia, including Afghanistan, Kohistan in northern Pakistan and Ladakh  
40 in northern India, though those two species in particular have not so far been recorded  
41 from the Lhasa Block of Tibet. Nevertheless, *S. tibetica* co-occurs with *Auroradiolites*  
42 *biconvexus* (Yang et al.), which probably evolved directly from *A. gilgitensis*  
43 (Douvillé), and the age of the latter association is in accordance with the generally

accepted age of the Yasin fauna as late Aptian to Albian. Hence the *S. tibetica* and *A. biconvexus* association can be considered a regional variant of the Yasin fauna, which had evidently already dispersed to the Lhasa Block by the late Aptian. So the Langshan Formation can be considered palaeogeographically linked with other mid-Cretaceous shallow-marine carbonate deposits in adjacent southwestern Asian regions. These findings also provide new evidence that the age of the rudist assemblage of the Lhasa Block is late Aptian to Albian, although a slightly younger age cannot be excluded.

**Keywords:** Lhasa Block; Langshan Formation; mid-Cretaceous; Rudists; Polyconitidae; Yasin fauna

## 1. Introduction

Rudists are an extinct order of bivalves that flourished on the carbonate platforms of the Tethyan–Atlantic–Pacific oceanic realm during the Late Jurassic to Cretaceous (Skelton, 2018). The mid-Cretaceous rudists of southwestern Asia and the western Pacific region show great similarity and affinity (Sano and Masse, 2013; Skelton et al., 2013). Accordingly, the Southwest Asian/Pacific Faunal Province was proposed, covering Iran, Afghanistan, Kohistan in northern Pakistan, Ladakh in northern India, Tibet in China and Japan, as well as Cebu Island and the Japanese seamounts in the Pacific region (Rao et al., 2015, 2017). The rudist fauna of this province is dominated by the endemic radiolitid *Auroradiolites* Rao et al., 2015 and the polyconitid lineage of *Horiopleura haydeni* Douvillé, 1926–*Praecaprotina* Yabe

and Nagao, 1926– *Magallanesia* Sano et al., 2014, as well as some more widely dispersed taxa such as *Eoradiolites* Douvillé, 1909 and *Sellaea* Di Stefano, 1889 (Skelton et al., 2013; Sano et al., 2014; Rao et al., 2015). The orbitolinid assemblage associated with these rudists is likewise composed of several endemic taxa such as *Mesorbitolina birmanica* (Sahni, 1937), *Palorbitolinoides orbiculata* Zhang, 1986 and *Palorbitolinoides hedini* Zhang, 1986, together with cosmopolitan taxa such as *Mesorbitolina texana* (Roemer, 1849), *M. subconcava* (Leymerie, 1878) and *M. aperta* (Erman, 1854) (Cherchi and Schroeder, 2013; Schlagintweit and Wilmsen, 2014; Rao et al., 2015, 2017; Boudagher–Fadel et al., 2017).

Within the SW Asian/Pacific Faunal Province, the so-called ‘Yasin fauna’ of rudists represented by the late Aptian–Albian *Horiopleura haydeni*/*Auroradiolites gilgitensis* association had a widespread distribution in southwest Asia including Iran, Afghanistan, Kohistan, and Ladakh (Douvillé, 1926; Rossi Ronchetti, 1965; Montenat et al., 1982; Upadhyay, 2001, 2014; Skelton et al., 2005; Sha and Cestari, 2016). In the mid-Cretaceous, these fossil sites were all located on blocks/terrane arrays along, or offshore from the southern Asian margin on the northern side of the eastern Neo-Tethyan Ocean (Rao et al., 2017). The Yasin fauna, as such, has not previously been recorded from the Lhasa Block, which occupied a similar geographical and tectonic position to the blocks/terrane arrays mentioned above. Instead, *Auroradiolites biconvexus*, probably derived directly from *A. gilgitensis*, and the canaliculate polyconitid genus *Magallanesia* were found there (Rao et al., 2015; 2017).

In this study, a new polyconitid genus, *Shajia*, which is superficially comparable

with *Horiopleura haydeni*, is described, based on articulated specimens collected from the Langshan Formation of Nyima County, Lhasa Block. A single specimen collected in Ladakh and formerly assigned to *Polyconites?* sp., by Masse and Fenerci-Masse (2017) is also reappraised in the light of the new material from the Lhasa Block. The relationship of *Shajia* with the Yasin fauna and its palaeogeographical implications are evaluated and the age of the rudist fauna from the Langshan Formation revised accordingly.

## 2. Geological setting

The Qinghai–Tibet plateau is generally regarded as a complex tectonic collage of several blocks; from north to south they are the Songpan–Ganze–Hoh Xil, Qiangtang, Lhasa and Tethyan–Himalayan blocks, which are separated from each other by sutures (Yin and Harrison, 2000; Zhu et al., 2013). These blocks rifted from Gondwana, drifted northward, and finally accreted to the Asian continent successively during the late Palaeozoic and Mesozoic eras, prior to the India–Asia collision (Pan et al., 2012; Li et al., 2017). Today, the Lhasa Block is bounded by the Bangong–Nu suture to the north and the Indus–Yarlung suture to the south (Fig. 1A; Yang et al., 2015). It rifted from the India–Gondwana continent in the Late Triassic, and collided with Asia in latest Jurassic–earliest Cretaceous times (Yin and Harrison, 2000; Metcalfe, 2006).

[Figure 1 hereabouts]

In the late Early to early Late Cretaceous (Barremian–Cenomanian), the Lhasa



Block had already collided with the Qiangtang Block, and was located on the north side of the eastern Tethys Ocean. At this time, shallow marine carbonate deposits, represented by the Langshan Formation, were widely distributed along the northern portion of this block (Leier et al., 2007; Rao et al., 2015). The Langshan Formation crops out as a nearly east-west-oriented belt extending from Baingoin County in the east to Rutog County in the west (XZBGM, 1993). It is dominated by dark-grey to black limestone sometimes interbedded with siltstone and mudstone (Zhang et al., 2004; Leier et al., 2007). The thickness of the Langshan Formation is poorly constrained and has considerable lateral variation; in this paper, we regard it as about 1000 m, referring to Zhang et al. (1986), Leeder et al. (1988), and Leier et al. (2007).

Abundant fossils have been recorded from the Langshan Formation, dominated by orbitolinids (Zhang, 1982, 1986, 1991; Boudagher-Fadel et al., 2017), rudists (Yang et al., 1982; Gou and Shi, 1998, Scott et al., 2010; Rao et al., 2015, 2017), gastropods (Yu and Xia, 1985) and corals (Löser and Liao, 2001; Deng and Wang, 2013). The age of the Langshan Formation is mainly constrained by foraminifers, especially large benthic orbitolinids. Deposition started from the Early Barremian based on the occurrence of *Eopalorbitolina charollaisi* Schroeder and Conrad, 1968 (Rao et al., 2015), and extended to the Early Cenomanian, as indicated by the appearance of *Daxia cenomana* Cuvillier and Szakall, 1949, *Nezzazata conica* (Smout, 1956), *Cuneolina* cf. *cylindrica*, *Orbitolina qatarica* Henson, 1948, *Conicorbitolina* sp. A and *Pseudedomia* sp. (Boudagher-Fadel et al., 2017).

The polyconitid specimens described herein were collected from the Langshan

Formation on the western side of Dangqiong Lake in southern Nyima County (Fig. 1B). This fossil site is located in the southern Nyima Basin, about 450 km northwest of Lhasa (DeCelles et al., 2007; Kapp et al., 2007). No palaeontological data have previously been published from the Langshan Formation of the Nyima Basin. According to the observations of Tong Zhang and Zhaoxiong Ma during field work, these specimens were collected from the lower part of the upper Langshan Formation. Furthermore, several large specimens of *Auroradiolites* were collected from the same bed as these polyconitid specimens. Although the left valves of these *Auroradiolites* specimens were not preserved, they are most likely attributable to *A. biconvexus* because of the large size of the right valves. As the age range of *A. biconvexus* was revised to late Aptian–Albian (Rao et al., 2017), and the geological range of the upper Langshan Formation is late Aptian to early Cenomanian (Rao et al., 2015), a late Aptian to Albian age can be assumed for these polyconitid specimens.

### 3. Systematic palaeontology

This published work and the nomenclatural acts it contains have been registered in Zoobank: <http://www.zoobank.org/References/xxxxxx>

The numbered specimens are housed at the Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, 39 East Beijing Road, Nanjing 210008, China (NIGPAS).

Abbreviations: *at*, anterior tooth; *ct*, central tooth; *Lam*, anterior myophore of left valve; *Lig*, Ligamentary infolding; *Lpm*, posterior myophore of left valve; *LV*, left valve; *ol*, outer shell layer; *pt*, posterior tooth; *Ram*, anterior myophore of right valve;

*Rpm*, posterior myophore of right valve; *RV*, right valve.

Quantitative characters (Fig. 2): *Dap*, antero–posterior commissural diameter, *Ddv*, dorso–ventral commissural diameter; *Hd*, dorsal height of *RV*; *Hv*, ventral height of *RV*; *LVc*, convexity of the *LV*.

The suprageneric classification used herein follows Skelton (2013a, b).

[Figure 2 hereabouts]

**Order** HIPPURITIDA Newell, 1965 (*nom. correct.* Scarlato & Starobogatov, 1971, *pro* Order Hippuritoida Newell, 1965; emend. Bouchet *et al.* 2010)

**Suborder** HIPPURITIDINA Newell, 1965 (Skelton, 2013b)

**Superfamily** RADIOLITOIDEA d’Orbigny, 1847

**Family** POLYCONITIDAE Mac Gillavry, 1937

**Genus** SHAJIA gen. nov.

*LSID*. urn:lsid:zoobank.org:act: xxxxxx

**Type species.** *Shajia tibetica* gen. et sp. nov.

**Derivation of name.** The new genus is named for Professor Jingeng Sha, a well-known Chinese palaeontologist who has made extensive studies of Mesozoic bivalves and of the geological evolution of the Qinghai–Xizang Plateau (Tibet). In recent years, he has focused on reviewing the rudist record of China, including publication of an important paper in *Cretaceous Research* to revise the late Aptian–Albian Yasin-type rudist fauna of the Himalayan area (Sha and Cestari, 2016). We chose his name as the root for the new genus erected herein in honour of his

contributions to Himalayan, and especially Tibetan rudist research.

**Diagnosis.** Large-sized polyconitid (antero-posterior commissural diameter can reach 90mm). *RV* conical and slightly twisted, *LV* low capuloid with dorsally over-hanging umbo. Ventral height of *RV* is less than twice the dorsal height, resulting in a dorsally inclined commissure. Outer shell layer is relatively thick, especially in the *RV* where its thickness may reach nearly 10 mm, on which longitudinal ribs are also developed. Subequal teeth of *LV* (*at* > *pt*) straddling the straight and erect central tooth of *RV*. Ligamentary infolding strongly developed. *LV* anterior myophore a narrow, flat-ended buttress projecting in from the anterior valve wall and facing onto the shelf-like anterior myophore of the *RV*, which is formed by a thickening of the inner shell layer. Posterior myophore in *LV* is a plate projected subparallel to the posterior valve wall, from which it is thus separated by a sub-rectangular ectomyophoral cavity; the adductor insertion surface on the broad tip of the *Lpm* is inclined somewhat outwards so as to face onto the depressed, inwardly sloping surface of the *RV* posterior myophore, which is formed by a pillow-like swelling of the inner shell layer.

[Figure 3 hereabouts]

**Discussion.** The thickening of the outer shell layer, especially of the *RV*, the subequal teeth of the *LV* (*at* > *pt*) and reflexed plate-like *Lpm* with adjacent ectomyophoral cavity, all observed in the Nyima specimens (Figs. 3A; 4-6) are diagnostic characters of the family Polyconitidae Mac Gillavry, 1937 (Skelton, 2013a). Comparisons thus need to be made with the handful of known genera in that family, starting with those

most easily excluded from further consideration.

The lack of sub-division into multiple canals of the *LV* posterior ectomyophoral cavity of the *Nyima* specimens (Figs. 3A) rules out the genus *Magallanesia* Sano et al., 2014 (= ‘Polyconitid new taxon 1’ in Skelton et al., 2013) (Fig. 3F). The genus *Polyconites* Roulland, 1830, can also be excluded as the *LV* of this genus is nearly flat (operculiform), the external surface of the shell lacks ribbing, and the posterior myophore of the *RV* is generally represented only by a slight thickening of the inner shell layer, at least in more derived species (Fig. 3D) (Masse et al., 1998; Skelton and Masse, 1998; Skelton, 2013a), although it is a little more prominent in juvenile shells of more primitive species (Skelton et al., 2010).

The external morphology of the *Nyima* specimens is similar to that of the genus *Horiopleura*, including the sparse, subdued longitudinal ribs (Fig. 4D), similar to those described from the Himalayan species *Horiopleura haydeni* by Douvillé (1926) and Rossi Ronchetti (1965). Moreover, the anterior myophoral arrangement of *Shajia* differs little from that seen in *Horiopleura* species (e.g., Fig. 3C). However, the depressed, inwardly sloping *Rpm* in the *Nyima* specimens (Fig. 3A) contrasts sharply with that in *Horiopleura haydeni*, which projects upward so as to face back onto the inner face of the *Lpm* (Fig. 3B). The same contrast is observed, moreover, in the type species of *Horiopleura*, *H. lamberti* Douvillé, 1889 (Fig. 3C), as well as in the more derived genera *Praecaprotina* (Fig. 3E), and ‘Polyconitid new taxa 2 and 3’ (Skelton et al., 2013), both of the latter additionally showing only a weakly convex *LV*.

Otherwise, both *Tepeyacia* and ‘Polyconitid new taxon 4’ (Skelton et al., 2013) are characterized by distinctive pleated infoldings of the *RV* outer shell layer, not observed in the Nyima specimens, while the polyconitid affinity of *Douvilleilia* and *Jerjesia* is in any case questionable (Skelton, 2013a; Masse et al., 2015) as neither appears to possess the diagnostic *LV* posterior ectomyophoral cavity.

In a recent review of the genus *Horiopleura*, however, Masse and Fenerci-Masse (2017, p. 54) noted that the outwardly inclined (facing towards the posterior side) and/or concave upward form of the *Rpm* (as in Fig. 3B, C) is ‘reported in the group of large, advanced species, e.g. *Horiopleura lamberti*; [whereas] in the group of small, putative primitive species, the inclination may be inwards (towards the anterior side), but a posterior shoulder is always present, a character that differentiates it from *Polyconites*’. All the figured specimens of the ‘putative primitive species’ assigned to *Horiopleura* by Masse and Fenerci-Masse (2017; e.g., Figs. 7C, 8E, F, 9A, B, 13A3 therein) indeed show this ‘shoulder’ to be a consistent attribute, such that adductor insertion area on the *Rpm* invariably forms a distinct ledge (either flat or slightly concave) that is oriented sub-parallel to the commissural plane.

Nevertheless, an Albian specimen from Parnassus, Greece, described by Masse and Fenerci-Masse (2017; Fig. 19B therein) and assigned by them to *H. distefanoi* Parona, 1909, does show an inwardly sloping *Rpm* with a correspondingly suppressed internal shoulder – a feature accordingly incorporated in their diagnosis for the species: ‘RV ledge like posterior myophore sloping inwards’ (Masse and Fenerci-Masse, 2017; p.

72). Accordingly, we suggest that the species that includes that specimen, if not *H. distefanoi* itself (pending confirmation the myophoral condition cited above in that species), might be allied with the new genus proposed herein, as an intermediate form, on the grounds that it differs from other species of *Horiopleura* by the marked inward inclination of the *Rpm* with associated suppression of the internal shoulder. This myophoral distinction is analogous to that which was used diagnostically to separate the earliest species of *Polyconites*, *P. hadriani*, from its inferred ancestral species of *Horiopleura* (Skelton et al., 2010), though additionally accompanied in that case by the pronounced flattening of the LV, in contrast to *Shajia*. Likewise, we propose that a single specimen from Ladakh that was assigned to *Polyconites?* sp., by Masse and Fenerci-Masse (2017), and which shows remarkably similar characteristics to those of the Nyima specimens (see below), should also be transferred to the genus *Shajia*.

**Age and distribution.** Late Aptian to Albian for the rudist-bearing limestone bed of Nyima County in Tibet (China), and late Aptian for the single specimen recorded from Ladakh (northern India); therefore, a late Aptian to Albian age should be assigned to this new genus based on current records.

*Shajia tibetica* sp. nov.

Figures 4–6

*LSID.* urn:lsid:zoobank.org:act: xxxxxx

? aff. 2017 *Horiopleura distefanoi* (Parona, 1909) Masse and Fenerci-Masse, fig. 19B

257 2017 *Polyconites* ? sp. Masse and Fenerci-Masse, fig. 20

258 **Derivation of name.** From Tibet, the type locality where the new species was found.

259 **Material.** Two articulated specimens (NIGP. 110522-110523), with the outer shell  
260 layer partially worn. The *RV* apices of both specimens were broken.

261 **Holotype.** Articulated specimen (NIGP. 110522; Figs. 4, 5). Two antero-posterior  
262 sections cutting through both valves were made (Fig. 4C), the more ventral section  
263 (Fig. 5A-B) oblique to the commissural plane and the more dorsal section (Fig. 5C-D)  
264 sub-perpendicular to it.

265 **Paratype.** Articulated specimen (NIGP. 110523; Fig. 6). Three obliquely transverse  
266 sections were made (Fig. 6A): the upper one cuts through both valves (the  
267 over-hanging dorsal part of the *LV* and the ventral part of the *RV*, Fig. 6C); the middle  
268 one cuts mainly across the *RV*, though including the umbonal tip of the *LV* (Fig. 6D,  
269 E); and the lower section cuts across the *RV* only (Fig. 6F).

270 **Localities.** The specimens were collected from the Langshan Formation of Nyima  
271 County, Tibet. The fossil site is located at the west side of Dangqiong Lake (E  
272  $86^{\circ}27'23''$ , N  $31^{\circ}36'53''$ ; Fig. 1B).

273 **Diagnosis.** As for genus.

274 [Figures 4–6 hereabouts]

275 **Description.**



**External morphology.** The two specimens are relatively large; although both *RV* apices are broken, values of *Hd* can be estimated at about 70 mm, and of *Hv* up to 120 mm, creating a dorsally inclined commissure. The *RV* is slightly twisted and curved and the *LV* is of low capuloid form, becoming gradually lower from the dorsal to ventral parts (Fig. 4C). The commissure is sub-rounded in the holotype, with a 90 mm diameter, whereas in the paratype it is elliptical because of dorso-ventral compression, with a 90 mm *Dap* and a 50 mm *Ddv*. Although the *ol* is partly worn or encrusted, blunt longitudinal ribs could be observed in each *RV* (Figs. 4D, 6B); rib width is about 2 mm. In the *LV*, fine costae radiate from the apex to the commissural rim.

The invaginated ligament, supported on an infolding of the dark outer shell layer, created a longitudinal furrow on the dorsal flank of the *RV* (Fig. 6A, F). The extent of external wear of the specimens precludes the possibility of recognizing radial bands either in the transverse sections, or on the natural surfaces of the shells.

**Internal organization.** The thickness of the dark outer shell layer may approach 10 mm in parts of the *RV* (Figs. 5, 6C-F). In the *LV*, in contrast, it is only a few mm thick, thickening slightly towards the commissure (Fig. 5). The inner shell layer is thick, especially in the myocardial area, and has been replaced by white to pale grey calcite spar (Figs. 5, 6 C, D, F). In the oblique transverse section of the paratype *RV*, the *ol* infolding associated with the invaginated ligament is about 3 mm wide and 6 mm in length, and a small ligamentary cavity wraps around its truncated inner tip (Fig. 6F).

The myocardial complex is well displayed in both specimens despite some localized boring and minor disruption by narrow, spar-filled fractures and stylolites.

The *Lpm* is prominent, pedunculate, and reflexed so as to lie nearly parallel to the over-arching posterior shell wall of the *LV*, from which it is thus separated by an ectomyophoral cavity of sub-rectangular shape in section (Fig. 5C, D); a narrow annexe extends dorsally from the latter cavity above the *pt* (Figs. 5A, 6C, 'o'). The tip of the *Lpm*, where the adductor muscle attached, faces outwards onto the depressed and inwardly sloping face of the *Rpm*, which is formed by a swelling of the *RV* inner shell layer (Fig. 5C, D). The *Rpm* thus appears in transverse section simply as a solid thickening of the posterior inner shell, lacking any indication of an ectomyophoral cavity behind it (Fig. 6D-F). The *Lam* is a protruding buttress with a bluntly sub-rectangular termination in section, separated from the anterior valve wall by a narrow gutter (Figure 5D). Its counterpart in the *RV*, the *Ram*, is formed by a shelf-like thickening of the inner shell layer (Figs. 5C, D, 6D-F). The *ct* in the *RV* is an erect, robust ridge interposed between the *at* and *pt* of the *LV* (Fig. 5A; 6D, E); *at* is slightly larger than *pt* (Fig. 6D-F).

**Remarks.** The single Ladakh specimen, which was assigned to *Polyconites* ? sp., by Masse and Fenerci-Masse (2017; Fig. 20 therein), is closely comparable with the Nyima specimens. It is similar to them in terms of the external morphology of the shell (including the low capuloid form of the *LV*), the longitudinal ribbing of the *RV*, and the myocardinal apparatus, with, in particular, a depressed, inward-sloping *Rpm* lacking a prominent internal shoulder (hence the appearance of 'strong affinity with *Polyconites*' noted by Masse and Fenerci-Masse, 2017, p. 72). Though somewhat smaller than the Nyima specimens, this specimen is virtually identical in form to the

type material found in Tibet and can thus be considered conspecific.

As noted earlier, in the discussion of the genus, the specimen assigned to *Horiopleura distefanoi* by Masse and Fenerci-Masse (2017) and thus perhaps that species – of similar age to *Shajia* – likewise shows a comparable myophoral configuration, as well as a *LV* of low capuloid shape. Taken together with other associated forms discussed by them, such similarities may bear upon the evolutionary origin of the new genus.

**Age and distribution.** As for genus.

## **4. Discussion**

### **4.1 Phylogenetic relationship of *Shajia* to other polyconitids**

At first sight, *Shajia* appears to have a chimaeric combination of characters from both *Horiopleura* and *Polyconites*: while its external shell form and ornamentation are very similar to those of the larger species of *Horiopleura*, the posterior myophoral arrangement, in particular, approaches that of *Polyconites*. Notwithstanding its distinction from both of those genera, derivation from either of them appears plausible: i.e., either from *Horiopleura*, through progressive inward inclination and depression of the *Rpm*, analogously to that proposed for the earlier origination of *P. hadriani* from an older *Horiopleura* sp. (Skelton et al., 2010), though without flattening of the *LV* and loss of ribbing; or, alternatively, from a *Polyconites* sp., through reversal of the *LV* flattening, to re-acquire a *Horiopleura*-like external shell shape, meanwhile retaining a depressed, inwardly-inclined *Rpm* – in contrast to the pronounced posteriorward tilt of the *Rpm* in the more derived species of *Horiopleura*.

Although *Shajia* is externally quite similar to the co-eval SW Asian *H. haydeni* in size, morphology, longitudinal ribs on the *RV*, as well as with respect to dentition, on the evidence of the few specimens known so far, it differs sharply from the latter species in its posterior myophoral organisation. Specifically, 1) the *Lpm* of *Shajia* is strongly reflexed posteriorly, such that it is nearly parallel to the posterior *LV* shell wall, forming a narrow sub-rectangular ectomyophoral cavity between them, whereas in *H. haydeni*, the *Lpm* protrudes more directly down into the *RV*; 2) correspondingly, the *Rpm* of *Shajia* is formed by the low, inward-facing swelling on the inner shell layer, whereas in *H. haydeni*, the *Rpm* is tilted backwards to form an erect plate with the adductor insertion surface on its posterior side (contrast Fig. 3A and B). As noted earlier, a similar contrast is seen also with respect to *H. lamberti*, as well as other, more derived polyconitid genera (Fig. 3 C, E, F).

On the other hand, co-eval species of *Polyconites* differ markedly from *Shajia* not only in their external form and ornamentation, as noted above, but also in the more extreme deflation of the *Rpm* (Fig. 3D). Furthermore, Sha and Cestari (2016) have drawn attention to the remarkable lack of records to date of *Polyconites* from the SW Asian Province (with the single exception of an unconfirmed mention of '*Polyconites* sp.', from Ladakh, by Mathur et al., 2008), weakening the circumstantial case for the derivation of *Shajia* from the latter genus.

Thus, by default, on the basis of the currently available evidence the most likely contender for the progenitor of *Shajia* is the species (or another, similar) represented by the specimen referred to *H. distefanoi* by Masse and Fenerci-Masse (2017; Fig.

19B therein), as discussed above. According to this scenario, *Shajia* would represent an iterative derivation of a *Polyconites*-like posterior myophoral organization from one of the ‘third group’ of *Horiopleura* species from the southern margin of the Mediterranean Tethys (extending to Oman) recognized by Masse and Fenerci-Masse (2017; pp. 75–76 therein). If correct, such a derivation would present an intriguing parallel with the appearance of another typically central to southern Tethyan form, *Sellaea* sp., in the Langshan fauna (Rao et al., 2015).

#### 4.2. The composition and age of the rudist fauna of the Langshan Formation

Study of rudists from the Langshan Formation was first conducted by Yang et al. (1982). Six species assigned to four genera were described from Rutog County, including *Toucasia* sp., *Requienia*? sp., *Rutonia bangonghuensis* Yang et al., 1982 and three species of *Praeradiolites*, *P. hedini* Douvillé, 1916, *P. biconvexus* Yang et al., 1982 and *P. ngariensis* Yang et al., 1982. Gou (1994) and Gou and Shi (1998) described eight new species plus an undetermined ninth, assigned to three genera, from the Langshan Formation of Coqen, Bangoin and Gegyai counties, comprising seven species placed in *Praeradiolites*, *P. gegyainensis* Gou, 1994, *P. exiguous* Gou, 1994, *P. perbellus* Gou, 1994, *P. gregareus* Gou, 1994, *P. daxungensis* Gou and Shi, 1998, *P. bangoinensis* Gou and Shi, 1998, *P. coquenensis* Gou and Shi, 1998, together with *Coralliochama anomalusa* Gou and Shi, 1998, and *Gyropleura*? sp.

Scott et al. (2010) restudied the specimens described by Yang et al. (1982) and considered the rudist assemblage of the Langshan Formation to be Albian to Cenomanian in age. They suggested that *Monopleura* sp., had been mistaken for

*Toucasia* sp., by Yang et al. (1982); they assigned *Praeradiolites ngariensis* to *Eoradiolites gilgitensis* based mainly on the structure of the anterior and posterior radial bands; and they attributed *P. biconvexus* to the genus *Sphaerulites* based on the external morphology.

Rao et al. (2015, 2017) revised the rudist taxa of the Langshan Formation described by Yang et al. (1982), Gou (1994), and Gou and Shi (1998). A new genus, *Auroradiolites*, was proposed for the endemic grouping of SW Asian to Japanese radiolitid species characterised by a compact *ol*, which were formerly attributed to *Praeradiolites* by Yang et al. (1982), Gou (1994), and Gou and Shi (1998). *Eoradiolites* cf. *hedini*, *Magallanesia rutogensis* and *Sellaea* sp. were also recognized, based on newly collected specimens. The age of the rudist assemblage reported by Rao et al. (2015, 2017) is most likely late Aptian to Albian, as the age range of *A. biconvexus* was revised to late Aptian to Albian by Rao et al. (2017).

As the age range of the Langshan Formation goes no younger than the early Cenomanian, according to orbitolinid data (Boudagher-Fadel et al., 2017), the rudist assemblage described from the Langshan Formation by Rao et al. (2015, 2017) is most probably late Aptian–Albian in age, though a younger age cannot be excluded.

### 4.3. Palaeobiogeographic implications

*Horiopleura haydeni*, and *Auroradiolites gilgitensis*, which together constitute the ‘Yasin-type’ rudist fauna (Skelton et al., 2005), were both first described by Douvillé (1926) as new species from the Yasin Group of the Gilgit region, northern Pakistan. This fossil site (Table1, No. 4) is located in the northern Kohistan terrane,

next to the North Suture Zone, which separates Kohistan from the Asian continent (Karakoram) to the north (Searle et al., 1999; Robertson and Collins, 2002; Khan et al., 2009). Subsequently, Pudsey et al. (1985) described rudists and orbitolinids from Kohistan, and proposed that the rudist-bearing limestone was of latest Aptian and/or early to middle Albian age.

[Table 1 hereabouts]

In the adjacent Ladakh terrane, *Horiopleura* sp., tentatively compared with *H. haydeni* by Upadhyay (2001), also co-occurs with *Auroradiolites gilgitensis*. This rudist assemblage was identified from limestones within the Saltoro Formation near the village of Shukur (Table1, No. 3) in the Nubra-Shyok valley region along the Shyok Suture Zone, which separates Ladakh from the Karakoram to the north (Upadhyay 2001, 2014). Upadhyay (2014) suggested a latest Aptian to early Albian age for this rudist assemblage.

Further west in Afghanistan, *Horiopleura haydeni* has been reported from two localities: Montenat et al. (1982) identified *H. haydeni* from the upper Aptian strata at the Adi Gar Mountains, about 30km north-northwest of Spin Boldak in the Kandahar region (Table1, No. 5); Rao et al. (2017) described two *H. haydeni* specimens, which were found together with *Auroradiolites gilgitensis*, from the Khist hills, south of Khash Rud, in central Afghanistan (Table1, No. 6).

*Horiopleura haydeni* thus has a relatively wide distribution extending along a nearly east-west-oriented belt in southwestern Asia, including central Afghanistan,

Kohistan in northern Pakistan and Ladakh in northern India (Fig. 7). These Yasin fauna species, *per se*, have not so far been recorded from the Lhasa Block, but instead *A. biconvexus* – probably derived from *A. gilgitensis* – and the canaliculate polyconitid genus *Magallanesia* were found there (Rao et al., 2015, 2017).

[Fig. 7 hereabouts]

*Shajia tibetica* was first recorded from Ladakh, as '*Polyconites* sp.', without detailed locality information, by Masse and Fenerci-Masse (2017). The specimens of *S. tibetica* recorded herein from the Langshan Formation of Nyima County, northern Lhasa Block (Fig. 7), were found in association with *A. biconvexus*, implying a late Aptian to Albian age (Rao et al., 2015), which is in broad agreement, moreover, with the generally accepted age of the Yasin fauna (Skelton et al., 2005; Sha and Cestari, 2016). Hence we conclude that *S. tibetica* and *A. biconvexus* can be regarded to be regional variants of the 'Yasin fauna'. Our material thus demonstrates that the Yasin fauna had already dispersed to the Lhasa Block by the late Aptian, which links the Langshan Formation palaeogeographically with other mid-Cretaceous sites of shallow-marine carbonate deposition in adjacent southwestern Asian regions.

The Kohistan–Ladakh terrane was situated between Asia to the north and India to the south in mid-Cretaceous times (Rolland et al., 2002). Although the exact timings of the collisions of the Kohistan–Ladakh terrane with Asia and with India, respectively, are still controversial (Rehman et al., 2011), it is generally accepted that the terrane itself was then located on the northern side of the Neo-Tethyan Ocean



(Chen et al., 1993; Searle et al., 1999). Meanwhile, according to the palaeogeographic reconstruction of Asia, central Afghanistan was situated on the southern Asian margin by mid-Cretaceous times (Chen et al., 1993; Zaman and Torii, 1999). At that time, the Lhasa Block had already collided with the Asian continent, and was also located at the northern margin of the Neo-Tethyan (Zhang, 2000; Leier et al., 2007). Therefore, all the localities that have yielded Yasin fauna were arrayed along or offshore from the southwestern Asian margin on the northern side of the Neo-Tethyan Ocean in the mid-Cretaceous.

As discussed above, in the SW Asian/Pacific Faunal Province, *Shajia* and *Horiopleura haydeni* were restricted to the southwestern Asia region. The genus *Praecaprotina* of late Aptian–Albian age has only been recorded from Japan and the Daiichi–Kashima Guyot in the Northwest Pacific (Masse and Shiba, 2010). The canaliculate polyconitid genus *Magallanesia*, of likely late Albian age, was found not only on Cebu Island and Takuyo–Daini Seamount, both located in the Pacific region (Sano et al., 2014), but also on the Lhasa Block in southwestern Asia (Rao et al., 2015). These polyconitid taxa, are all endemic to the combined southwestern Asia and Pacific regions, and mostly co-occur with the radiolitid genus *Auroradiolites*, which is characterised by an entirely compact outer shell layer and is also restricted to these regions (Rao et al., 2017). Hence these findings suggest strong endemism in the rudist fauna of the SW Asian/Pacific Faunal Province during the late Aptian to Albian interval.

## 5. Conclusions

*Shajia tibetica* gen. et sp. nov., a new polyconitid rudist, is described from the Langshan Formation of Nyima County in the northern Lhasa Block of Tibet. It is similar in external form to *Horiopleura haydeni* Douvillé, 1926, an abundant species of late Aptian to Albian age that is endemic to southwestern Asia, including Afghanistan, Kohistan in northern Pakistan, and Ladakh in northern India. However, *Shajia* differs from the latter species in its posterior myophoral arrangement: 1) *Lpm* is strongly bent posteriorly, such that it is nearly parallel to the over-arching posterior shell wall of the *LV*, thereby forming a sub-rectangular ectomyophoral cavity between them; 2) *Rpm* is formed by an inward-inclined swelling of the inner shell layer, without forming a projecting plate. A single specimen from Ladakh which was assigned to *Polyconites?* sp. by Masse and Fenerci-Masse (2017), is reappraised and transferred to *Shajia*. It was most likely derived from one of a small group of *Horiopleura* species that dwelt on the southern margin of the Mediterranean Tethys (extending to Oman) and which show a posterior myophoral arrangement approaching that of *Shajia*. The geological range of *Shajia* is probably Upper Aptian to Albian, combining the records from the Lhasa Block and Ladakh.

*S. tibetica* is associated with *A. biconvexus*, and as the age of the association is basically in accordance with that of the Yasin fauna (*H. haydeni* and *A. gilgitensis*) it may thus be considered as a regional variant of the latter association. This new material thus shows that the Yasin fauna had already dispersed to the Lhasa Block by the late Aptian, making the Langshan Formation comparable palaeobiogeographically with other mid-Cretaceous shallow-water carbonate deposits in adjacent southwestern

493 Asia regions.

## 494 **Acknowledgements**

495 This work was supported by the National Natural Science Foundation of China  
 496 [grant numbers 41702014, 41730317 and 41302011], the State Key Laboratory of  
 497 Palaeobiology and Stratigraphy [grant number 20162110], the Strategic Priority  
 498 Research Program of the Chinese Academy of Sciences [grant number  
 499 XDB26000000], and by the China Geological Survey [grant number DD20160015]. It  
 500 is also a contribution to UNESCO–IUGS IGCP (United Nations Educational,  
 501 Scientific and Cultural Organization–International Union of Geological Sciences)  
 502 Project 632. We warmly thank our referees, Jean–Pierre Masse and Eulàlia Gili for  
 503 their detailed and constructive critical comments, which greatly helped us to improve  
 504 the presentation and arguments of our paper.

## 505 **References**

- 506 BOUCHET, P., ROCROI, J.—P., BIELER, R., CARTER, J.G. and COAN, E.V., 2010.  
 507 Nomenclator of Bivalve Families (P. B. and J.—P. R.) and Classification of Bivalve  
 508 Families (R.B., J.G.C. and E.V.C.). *Malacologia*, **52**, 1–184.
- 509 BOUDAGHER—FADEL, M. K., HU, X., PRICE, G. D., SUN, G., WANG, J. and AN,  
 510 W., 2017. Foraminiferal Biostratigraphy and Palaeoenvironmental Analysis of the  
 511 Mid--Cretaceous Limestones in the Southern Tibetan Plateau. *Journal of*  
 512 *Foraminiferal Research.*, **47(2)**, 188–207. (in press).
- 513 CHEN, Y., COURTILLOT, V., COGNÉ, J. P., BESSE, J., YANG, Z. and ENKIN, R.,

1993. The Configuration of Asia Prior to the Collision of India: Cretaceous  
Paleomagnetic Constrains. *Journal of Geophysical Research*, **98**, 927–941.
- CHERCHI, A. and SCHROEDER, R., 2013. The *Praeorbitolina/Palorbitolinoides*  
Association: an Aptian Biostratigraphic Key—Interval at the Southern Margin of  
the Neo—Tethys. *Cretaceous Research*, **39**, 70–77.
- CUVILLIER, J. AND SZAKALL, V., 1949. Foraminifères d'Aquitaine, Première  
partie (Reophacidae à Nonionidae), Boisseau Imp., Toulouse, 112 pp.
- D'ORBIGNY, A. D., 1847. Sur les Brachiopodes ou Palliobranches (deuxième  
Mémoire). *Comptes Rendus Hebdomadaires des Séances de l'Académie des*  
*Sciences (Paris)*, **25**, 266–269.
- DECELLES, P. G., KAPP, P., DING, L. and GEHRELS, G. E., 2007. Late  
Cretaceous to middle Tertiary basin evolution in the central Tibetan Plateau:  
Changing environments in response to tectonic partitioning, aridification, and  
regional elevation gain. *Geological Society of America Bulletin*, **119** (5–6), 654–  
680.
- DENG, Z. Q. and WANG, G. X., 2013. Mesozoic Corals and Sponges from Gê'gyai  
and Rutog, Xizang. *Acta Palaeontologica Sinica*, **52**, 202–212. [In Chinese with  
English abstract]
- DI STEFANO, G., 1889. Studi stratigrafici e paleontologici sul sistema Cretaceo della  
Sicilia. 1. Gli strati con *Caprotina* di Termini—Imerese. *Atti della Reale Accademia*  
*di Scienze, Lettere e Belle Arte di Palermo*, Nuova Serie, **10**, 1–44.
- DOUVILLÉ, H., 1889. Sur quelques rudistes du terrain crétacé inférieur des Pyrénées.

- 536 *Bulletin de la Société géologique de France*, (3), **17**, 627–635.
- 537 DOUVILLÉ, H., 1909. Sur le genre *Eoradiolites* nov. *Comptes Rendus des Séances*  
538 *de la Société Géologique de France* (série 4), **9**, 77.
- 539 DOUVILLÉ, H., 1916. Les calcaires a Orbitolines et a Radiolites du Thibet. 145–147.  
540 In HEDIN, S. (ed.) *Southern Tibet – Discoveries in former times compared with my*  
541 *own researches in 1906–1908. Vol. 5 Petrographie und Geologie I. Beschreibung*  
542 *der Gesteinsproben, Anhang. Lithographic Institute of the General Staff of the*  
543 *Swedish Army, Stockholm.*
- 544 DOUVILLÉ, H., 1926. Fossiles recueillis par Hayden dans le Kashmir en 1906 et les  
545 Pamirs en 1914; leur description. *Records of the Geological Survey of India*, **58**,  
546 349–357.
- 547 ERMAN, A., 1854. Einige Beobachtungen über die Kreideformation an der  
548 Nordküste von Spanien. *Zeitschrift der Deutschen. Geologischen Gesellschaft*, **6**,  
549 596–611.
- 550 GOU, Z. H. and SHI, H., 1998. Rudists (Bivalvia) from the Cretaceous of Tibet,  
551 China, with Descriptions of New Species. 255–266. In JOHNSON, P.A. and  
552 HAGGART, J.W. (eds.). *Bivalves: an eon of evolution – paleobiological studies*  
553 *honoring Norman D. Newell. University of Calgary Press, Calgary, Canada, 461*  
554 *pp.*
- 555 GOU, Z. H., 1994. Rudists Fossils of Cretaceous Period in Gegyai Region, Tibet.  
556 *Tibet Geology*, **12**, 128–138. [In Chinese with English abstract]
- 557 HAY, W.W., DeCONTO, R., WOLD, C.N., WILSON, K.M., VOIGHT, S., SCHULZ,

- 558 M., WOLD—ROSSBY, A., DULLO, W.—C., RONO, A.B., BALUKHOVSKY,  
 559 A.N. and SOEDING E., 1999. Alternative Global Cretaceous Palaeogeography. In  
 560 BARRERA, E. and JOHNSON, C. (eds.). *The Evolution of Cretaceous*  
 561 *Ocean/Climate Systems, Geological Society of America Special Paper*, **332**, 1–47.
- 562 HENSON, F. R. S., 1948. Larger imperforate foraminifera of south-western Asia.  
 563 Families Lituolidae, Orbitolinidae and Meandropsinidae. *Monograph British*  
 564 *Museum (Natural History)*, London, 127pp.
- 565 KAPP, P., DECELLES, P. G., GEHRELS, G. E., HEIZLER, M. AND DING, L.,  
 566 2007. Geological records of the Lhasa—Qiangtang and Indo—Asian collisions in  
 567 the Nima area of central Tibet. *Geological Society of America Bulletin*, **119** (7–8),  
 568 917–933.
- 569 KHAN, S. D., WALKER, D. J., HALL, S. A., BURKE, K. C., SHAH, M. T. and  
 570 STOCKLI, L., 2009. Did the Kohistan—Ladakh Island Arc Collide First with India?  
 571 *Geological Society of America Bulletin*, **121**, 366–384.
- 572 LEEDER, M. R., SMITH, A. B., and JIXIANG, Y., 1988. Sedimentology,  
 573 Palaeoecology and Palaeoenvironmental Evolution of the 1985 Lhasa to Golmud  
 574 Geotraverse. *Royal Society of London, Philosophical Transactions, Series A*, **327**,  
 575 107–143.
- 576 LEIER, A. L., DECELLES, P. G., KAPP, P. and GEHRELS, G. E., 2007. Lower  
 577 Cretaceous Strata in the Lhasa Terrane, Tibet, with Implications for Understanding  
 578 the Early Tectonic History of the Tibetan Plateau. *Journal of Sedimentary Research*,  
 579 **77**, 809–825.

- 580 LEYMERIE, A., 1878, 1881. Description Géologique et Paléontologique des  
581 Pyrénées de la Haute-Garonne. E. Privat, Toulouse, 2 v. [text (1878): 1010 p. atlas  
582 (1881)].
- 583 LI, Z., DING, L., SONG, P., FU, J. and YUE, Y., 2017. Paleomagnetic Constraints on  
584 the Paleolatitude of the Lhasa Block during the Early Cretaceous: Implications for  
585 the Onset of India – Asia Collision and Latitudinal Shortening Estimates Across  
586 Tibet and Stable Asia. *Gondwana Research*, **41**, 352–372.
- 587 LÖSER, H. and LIAO, W. H., 2001. Cretaceous Corals from Tibet (China) --  
588 Stratigraphic and Palaeobiogeographic Aspects. *Journal of Asian Earth Sciences*,  
589 **19**, 661–667.
- 590 MAC GILLAVRY, H. J., 1937. Geology of the Province of Camagüey, Cuba, with  
591 Revisional Studies in Rudist Paleontology. *Geographische en Geologische*  
592 *Mededeelingen, Physiographisch--Geologische Reeks*, **14**, 168 pp.
- 593 MASSE, J. P. and FENERCI-MASSE, M., 2017. Taxonomy and Stratigraphy of Late  
594 Barremian–Albian Species of *Horiopleura* Douvillé (Hippuritida, Polyconitidae) of  
595 the Mediterranean and Southwestern Asian Regions. *Cretaceous Research*, **76**, 53–  
596 80.
- 597 MASSE, J. P., ARIAS, C. and VILAS, L., 1998. Lower Cretaceous Rudist Faunas of  
598 Southeast Spain: an Overview. In: MASSE, J. P., SKELTON, P.W. (Eds.),  
599 *Quatrième Congrès International sur les Rudistes*, Géobios, Mémoire Spécial, **22**,  
600 193–210.
- 601 MASSE, J. P., MOURGES, F. A. and FENERCI-MASSE. M., 2015. Aptian–Albian

- 602 Rudist Bivalves (Hippuritida) from the Chilean Central Andes: Their  
 603 Palaeoceanographic Significance. *Cretaceous Research*, **54**, 243–254.
- 604 MASSE, J.P. and SHIBA, M., 2010. *Praecaprotina kashimae* nov. sp. (Bivalvia,  
 605 Hippuritacea) from the Daiichi–Kashima Seamount (Japan Trench). *Cretaceous*  
 606 *Research*, **31**(2), 147–153.
- 607 MATHUR, N.S., JUYAL, K.P. and KUMAR, K. 2008. Biotic response to  
 608 Cretaceous–Eocene tectonic events at the northern margin of the Indian Plate and  
 609 the Indus–Tsangpo Suture Zone, Ladakh Himalaya, India. *Journal of the*  
 610 *Palaeontological Society of India*, **53**, 37–49.
- 611 METCALFE, I., 2006. Palaeozoic and Mesozoic Tectonic Evolution and  
 612 Palaeogeography of East Asian Crustal Fragments: The Korean Peninsula in  
 613 Context. *Gondwana Research*, **9**, 24–46.
- 614 MONTENAT, C., MOULLADE, M. and PHILIP, J., 1982. Le Crétacé inférieur à  
 615 Orbitolines et Rudistes d'Afghanistan central. *Géologie Méditerranéenne*, **9**, 109–  
 616 122.
- 617 NEWELL, N. D., 1965. Classification of the Bivalvia. *American Museum Novitates*,  
 618 **2206**, 1–25.
- 619 PAN, G., WANG, L., LI, R., YUAN, S., JI, W., YIN, F., ZHANG, W. and WANG,  
 620 B., 2012. Tectonic Evolution of the Qinghai–Tibet Plateau. *Journal of Asian Earth*  
 621 *Sciences*, **53**, 3–14.
- 622 PARONA, C.F., 1909. La fauna coralligena del Cretaceo dei Monti d'Ocre  
 623 nell'Abruzzo aquilano. *Memoria per servire alla descrizione della carta geologica*



- 624 *d'Italia*, **5**, 1–233.
- 625 PUDSEY, C. J., SCHROEDER, R. and SKELTON, P. W., 1985. Cretaceous
- 626 (Aptian/Albian) Age for Island–Arc Volcanics, Kohistan, N. Pakistan. 150–168. *In*
- 627 GUPTA, V. J. (ed.). *Contributions to Himalayan Geology, v. 3, Geology of western*
- 628 *Himalayas*. Hindustan Publishing Corporation, Delhi.
- 629 RAO, X., SKELTON P. W., SANO S., LI, C., PAN, Y., LUO, H., CAI, H., PENG, B.
- 630 and SHA, J., 2017. Evolution and Palaeogeographical Dispersion of the Radiolitid
- 631 Rudist genus *Auroradiolites* (Bivalvia: Hippuritida), with Descriptions of New
- 632 Material from Tibet and Archived Specimens from Afghanistan, *Papers in*
- 633 *Palaeontology*, **3(2)**, DOI: 10.1002/spp2.1076.
- 634 RAO, X., SKELTON, P.W., SHA, J., CAI, H. and IBA, Y., 2015. Mid–Cretaceous
- 635 Rudists (Bivalvia: Hippuritida) from the Langshan Formation, Lhasa block, Tibet.
- 636 *Papers in Palaeontology*, **1**, 401–424.
- 637 REHMAN, H. U., SENO, T., YAMAMOTO, H. and KHAN, T., 2011. Timing of
- 638 Collision of the Kohistan–Ladakh Arc with India and Asia: Debate. *Island Arc*, **20**,
- 639 308–328.
- 640 ROBERTSON, A. H. F. and COLLINS, A. S., 2002. Shyok Suture Zone, N Pakistan:
- 641 Late Mesozoic–Tertiary Evolution of a Critical Suture Separating the Oceanic
- 642 Ladakh Arc from the Asian Continental Margin. *Journal of Asian Earth Science*, **20**,
- 643 309–351.
- 644 ROEMER, F., 1849. *Texas*. Bonn, 464 pp.
- 645 ROLLAND, Y., PICARD, C., PÊCHER, A., LAPIERRE, H., BOSCH, D. and

- 646 KELLER, F., 2002. The Ladakh Arc of NW Himalaya: Slab Melting and Mantle  
647 Interaction During Fast Northward Drift of Indian Plate. *Chemical Geology*, **182**,  
648 139–178.
- 649 ROSSI RONCHETTI, C., 1965. Rudiste e nerinee del cretaceo di Yasin (Pakistan  
650 nord–occidentale). In DESIO A. (leader), BRILL E. J. (eds) *Italian Expeditions to*  
651 *the Karakorum (K2) and Hindu Kush, Scientific Reports*. Part 4, 229–272. Leiden.
- 652 ROULLAND, F., 1830. Nouvelles observations sur les *Ichthyosarcolites*. Actes de la  
653 Société Linéenne de Bordeaux, **4**, 164–166.
- 654 SAHNI, M. R., 1937. Discovery of *Orbitolina*-bearing Rocks in Burma, with a  
655 Description of *Orbitolina birmanica* sp. nov. *Records of the Geological Survey of*  
656 *India*, **71** (4), 360–375.
- 657 SANO, S. and MASSE, J. P., 2013. First record of a primitive radiolitid rudist from  
658 Japan. *Paleontological Research*, **17**, 317–324.
- 659 SANO, S., IBA, Y., SKELTON, P. W., MASSE, J.—P. AGUILAR, Y. M. and KASE,  
660 T., 2014. The Evolution of Canaliculate Rudists in the Light of a New Canaliculate  
661 Polyconitid Rudist from the Albian of the Central Pacific. *Palaeontology*, **57**, 951–  
662 962.
- 663 SCARLATO, O. A. and STAROBOGATOV, Y. I., 1971. Osnovnye Cherty Evolyutsii  
664 i sistema Klassa Bivalvia Morfologiya, Sistematika i Filogeniya Mollyuskov. *Trudy*  
665 *Zoologicheskogo Instituta, Akademiya Nauk SSSR*, **80**, 5–38. [In Russian]
- 666 SCHLAGINTWEIT, F. and WILMSEN, M., 2014. Orbitolinid Biostratigraphy of the  
667 Top Taft Formation (Lower Cretaceous of the Yazd Block, Central Iran).

- 668 *Cretaceous Research*, **49**, 125–133.
- 669 SCHROEDER, R. and CONRAD, M. A., 1968. Huitième note sur les Foraminifères  
670 du Crétacé inférieur de la région genevoise. *Eopalarbitolina charollaisi*, n. gen., n.  
671 sp., un Orbitolinidé nouveau du Barrémien à faciès urgonien. Compte rendu des  
672 séances de la Société de Physique et d'Histoire Naturelle de Genève, nouvelle série,  
673 **2(3)**, 145–162.
- 674 SCOTT, R. W., WAN, X., SHA, J. and WEN, S., 2010. Rudists of Tibet and Tarim  
675 Basin, China: Significance to Requieriidae Phylogeny. *Journal of Paleontology*, **84**,  
676 444–465.
- 677 SEARLE, M. P., KHAN, M. A., FRASER, J. E., GOUGH, S. J. and JAN, M. Q., 1999.  
678 The Tectonic Evolution of the Kohistan Collision Belt Along the Karakoram  
679 Highway Transect, North Pakistan. *Tectonics*, **18**, 929–949.
- 680 SHA, J. and CESTARI, R., 2016. Late Aptian–Albian Yasin-type Rudist  
681 Assemblage in the Himalayas: Palaeobiogeographic Implications. *Cretaceous*  
682 *Research*, **65**, 34–47.
- 683 SKELTON, P. W. and MASSE, J.—P., 1998. Revision of the Lower Cretaceous Rudist  
684 Genera *Pachytraga* Paquier and *Retha* Cox (Bivalvia: Hippuritacea), and the  
685 origins of the Caprinidae. *Géobios, mémoire spécial*, **22**, 331–370.
- 686 SKELTON, P. W., 2013a. Rudist Classification for the Revised Bivalvia Volumes of  
687 the 'Treatise on Invertebrate Paleontology'. *Caribbean Journal of Earth Science*,  
688 **45**, 9–33.
- 689 SKELTON, P. W., 2013b. Rudist Classification: Nomenclatural Correction of

- 690 'Suborder Radiolitidina Skelton, 2013' to 'Suborder Hippuritidina Newell, 1965'.  
 691 *Caribbean Journal of Earth Science*, **45**, 34.
- 692 SKELTON, P. W., 2018. Part N, Volume 1, Chapter 26A: Introduction to the  
 693 Hippuritida (rudists): Shell structure, anatomy, and evolution. *Treatise Online*  
 694 104:1–37.
- 695 SKELTON, P. W., SANO, S. and MASSE, J.—P., 2013. Rudist Bivalves and the  
 696 Pacific in the Late Jurassic and Early Cretaceous. *Journal of the Geological Society*,  
 697 **170**, 513–526.
- 698 SKELTON, P.W. and MASSE, J. P., 2000. Synoptic Guide to the Lower Cretaceous  
 699 Rudist Bivalves of Arabia. *SEPM Special Publication*, **69**, 85–95.
- 700 SKELTON, P.W. and SMITH, A.D., 2000. A Preliminary Phylogeny for Rudist  
 701 Bivalves: Sifting Clades from Grades. In: HARPER, E.M., TAYLOR, J.D.,  
 702 CRAME, J.A. (Eds.), *the Evolutionary Biology of the Bivalves*. Geological Society,  
 703 London. Special Publication, **177**, 97–127.
- 704 SKELTON, P.W., GILI, E., BOVER—ARNAL, T., SALAS, R. and MORENO—  
 705 BEDMAR, J.A., 2010. A New Species of *Polyconites* from the Lower Aptian of  
 706 Iberia and the Early Evolution of Polyconitid Rudists. *Turkish Journal of Earth*  
 707 *Sciences*, **19**, 557–572.
- 708 SKELTON, P.W., RAISOSSADAT, N., UPADHYAY, R. and BERNOULLI, D.  
 709 2005. 'Yasin--type' rudist fauna from eastern Iran and northern Ladakh. In  
 710 FILKORN, H.N. (Ed.), *SEPM Abstracts and Post—congress Field Guide*, Seventh  
 711 International Congress on Rudists, p. 91. Austin. ISBN 1565761197.

- 712 SMOUT, A. H., 1956. Three new Cretaceous genera of foraminifera related to the  
713 Ceratobuliminidae. *Micropaleontology*, **2**, 335–348.
- 714 STEUBER, T., 1999. Cretaceous Rudists of Boeotia, Central Greece. *Special Papers*  
715 *in Palaeontology*, **61**, 229 pp.
- 716 UPADHYAY, R., 2001. Middle Cretaceous Carbonate Build—Ups and Volcanic  
717 Seamount in the Shyok Suture, Northern Ladakh, India. *Current Science*, **81**, 695–  
718 699.
- 719 UPADHYAY, R., 2014. Palaeogeographic Significance of 'Yasin--type' Rudist and  
720 Orbitolinid Fauna of the Shyok Suture Zone, Saltoro Hills, Northern Ladakh, India.  
721 *Current Science*, **106**, 223–228.
- 722 WANG, L., PAN, G., DING, J., and YAO, D., 2013. *Geological Mapping illumination*  
723 *(1:1500000) of Qinghai–Tibet Plateau and Its adjacent area*. Geological Publishing  
724 House, Beijing. 288pp. [In Chinese]
- 725 XZBGM (XIZANG BUREAU OF GEOLOGY and MINERAL RESOURCES), 1993.  
726 *Regional geology of Xizang Autonomous Region*. Geological Publishing House,  
727 Beijing. 707pp. [In Chinese with English summary]
- 728 YABE, H. and NAGAO, S., 1926. *Praecaprotina*, nov. gen. from the Lower  
729 Cretaceous of Japan. *Science Reports of the Tohoku Imperial University, Second*  
730 *Series*, **9**, 21–24.
- 731 YANG, T., MA, Y., ZHANG, S., BIAN, W., YANG, Z., WU, H., LI, H., CHEN, W.  
732 and DING, J., 2015. New Insights into the India – Asia Collision Process from  
733 Cretaceous Paleomagnetic and Geochronologic Results in the Lhasa Terrane.

- 734 *Gondwana Research*, **28**, 625–641.
- 735 YANG, Z., NIE, Z., WU, S. and LIANG, D., 1982. Cretaceous Rudists from Ngari,  
 736 Xizang (Tibet), Autonomous Region, China and Their Geologic Significance. *Acta*  
 737 *Geologica Sinica*, **56**, 293–301. [In Chinese with English abstract]
- 738 YEHARA, S., 1920. A Pachyodont Lamellibranchia from the Cretaceous Deposits of  
 739 Miyako in Rikuchu. *Journal of the Geological Society of Tokyo*, **27**, 39–44.
- 740 YIN, A. and HARRISON, T. M., 2000. Geologic Evolution of the Himalayan—  
 741 Tibetan Orogen. *Annual Review of Earth and Planetary Sciences*, **28**, 2211–2280.
- 742 YU, W. and XIA, J. B., 1985. Late Jurassic–Cretaceous Nerineids from Northern  
 743 Xizang. *Acta Palaeontologica Sinica*, **24**, 640–650. [In Chinese with English  
 744 abstract].
- 745 ZAMAN, H. and TORII, M., 1999. Paleomagnetic Study of Cretaceous Red Beds  
 746 from the Eastern Hindukush Ranges, Northern Pakistan; Paleogeographic Construction of  
 747 the Kohistan–Karakoram Composite Unit before the India–Asia Collision.  
 748 *Geophysical Journal International*, **136**, 719–738.
- 749 ZHANG, B. G., 1982. *Orbitolina* (Foraminifera) from Xizang. 51–80. *In The Series*  
 750 *of the Scientific Expedition to the Qinghai–Xizang (Tibet) Plateau, Paleontology of*  
 751 *Xizang (Tibet)*, 4. Science Press, Beijing. [In Chinese with English abstract].
- 752 ZHANG, B. G., 1986. Early Cretaceous Orbitolinids from Xainza and Baingoin,  
 753 Xizang. *In Bulletin of Nanjing Institute of Geology and Palaeontology, Academia*  
 754 *Sinica*, **10**, 101–122. [In Chinese with English abstract].
- 755 ZHANG, B. G., 1991. Cretaceous Larger Foraminifera Orbitolines from Ngari,

Western Xizang (Tibet). 68–87. In SUN, D. L., XU, J. T. et al. (eds). *Stratigraphy and Palaeontology of Permian, Jurassic and Cretaceous from the Rutog Region, Xizang (Tibet)*. Nanjing University Press, Nanjing. [In Chinese with English abstract].

ZHANG, B. G., YANG, S. Q., SUN, D. L. and YE, C. H., 1986. Cretaceous System. In *Bulletin of Nanjing Institute of Geology and Palaeontology, Academia Sinica*, **10**, 46–55. [In Chinese with English abstract].

ZHANG, K. J., 2000. Cretaceous Palaeogeography of Tibet and Adjacent Areas (China): Tectonic Implications. *Cretaceous Research*, **21**, 23–33.

ZHANG, K. J., XIA, B. D., WANG, G. M., LI, Y. T. and YE, H. F., 2004. Early Cretaceous Stratigraphy, Depositional Environments, Sandstone Provenance, and Tectonic Setting of Central Tibet, Western China. *Geological Society of America, Bulletin*, **116**, 1202–1222.

ZHU, D., ZHAO, Z., NIU, Y., DILEK, Y., HOU, Z. and MO, X., 2013. The Origin and Pre–Cenozoic Evolution of the Tibetan Plateau. *Gondwana Research*, **23**, 1429–1459.

## Table caption

Table 1. Localities yielding *Shajia* and *Horiopleura haydeni* in the SW Asian/Pacific Faunal Province, with currently assigned ages. See Fig. 7 for the geographical distribution of these localities.

## Figure Captions

1. Location of study area. (A) Map showing the tectonic framework of Tibet and the outcrops of the Langshan Formation (grey) in the Lhasa Block (after XZBGM, 1993; Wang et al., 2013). (B) Locality map for *Shajia* in Nyima County (red rectangle in A). *Abbreviations*: BNS, Bangong–Nu Suture; IYS, Indus–Yarlung Suture.

[in colour]

2. Abbreviations of quantitative characters used for the description of *Shajia* (*LV*, left valve; *RV*, right valve): *Dap*, antero-posterior commissural diameter, *Ddv*, dorso–ventral commissural diameter; *Hd*, dorsal height of *RV*; *Hv*, ventral height of *RV*; *LVC*, convexity of the *LV*.

[in black and white]

3. Diagrammatic antero-posterior sections across both valves of *Shajia* and other similar polyconitids (outer shell layer shown in black; inner shell, in grey). A, *Shajia* (based on Fig. 5D herein); B, *Horiopleura haydeni* (after Rossi Ronchetti, 1965, Plate 38, fig. 1; outer shell layer worn, and image reversed to facilitate comparison); C, *Horiopleura lamberti* (after Skelton and Smith, 2000, Fig. 7); D, *Polyconites* (after Skelton, 2013a, Fig. 7A); E, *Praecaprotina yaegashii* (after



Sano et al., 2014, Fig. 2); F, *Magallanesia canaliculata* (after Sano et al., 2014, Fig. 4D). Abbreviations: *Lam*, anterior myophore of left valve; *Lpm*, posterior myophore of left valve; *LV*, left valve; *Ram*, anterior myophore of right valve; *Rpm*, posterior myophore of right valve; *RV*, right valve.

[in black and white]

4. *Shajia tibetica* gen. et sp. nov., holotype articulated specimen (NIGP. 110522). A, dorsal view showing overhanging umbo of *LV*. B, ventral view. C, anterior view showing the oblique commissure and the convexity of the *LV*; red lines show the two sections made on this specimen; a, b, c, and d represent the positions of the sections shown in Figs. 5A–D respectively. D, posterior view of *RV* showing the longitudinal ribs (inset: magnified view of red rectangle).

[in colour]

5. *Shajia tibetica* gen. et sp. nov., holotype (NIGP. 110522): A–D, successively more ventrally situated antero-posterior sections of both valves, as indicated either side of the red lines in Fig. 4C (B and D are reversed so that all images are viewed looking dorsally, for ease of comparison): A–B show the dentition; C–D show the myophoral organisation. Abbreviations: *at*, anterior tooth (from *LV*); *ct*, central tooth (from *RV*); *Lam*, anterior myophore of left valve; *Lpm*, posterior myophore of left valve; *LV*, left valve; *o*, annexe extending dorsally from the ectomyophoral cavity; *pt*, posterior tooth (from *LV*); *Ram*, anterior myophore of right valve; *Rpm*, posterior myophore of right valve; *RV*, right valve.

[in colour]

6. *Shajia tibetica* gen. et sp. nov., paratype articulated specimen (NIGP. 110523). A, dorsal view, commissure indicated by dashed red line; solid red lines show the three transverse sections made of this specimen. B, anterior view showing the longitudinal ribs on the RV. C, D–E and F show the three successively lower transverse sections indicated in A, all in adumbonal view of the RV, revealing the myocardinal organisation.

Abbreviations: *at*, anterior tooth (from LV); *ct*, central tooth (from RV); *Lig*, infolding of outer shell layer associated with invaginated ligament; *Lpm*, posterior myophore of left valve (dorsalmost part only); *LV*, left valve; *o*, annexe extending dorsally from the ectomyophoral cavity; *ol*, outer shell layer (of RV); *pt*, posterior tooth (from LV); *Ram*, anterior myophore of right valve; *Rpm*, posterior myophore of right valve; *RV*, right valve.



[in colour]


7. Present-day geographical map showing the fossil localities bearing *Shajia* and *Horiopleura haydeni*, represented by red and blue stars respectively (see Table 1 for details). The base map was generated using the ODSN online Plate Tectonic Reconstruction Service (Hay et al., 1999). CAF, central Afghanistan; KOH–LAD, Kohistan–Ladakh terrane; LSA, Lhasa block.


[in colour]

Table 1


Record	Locality	Age	Species	References
1	Nyima County, Tibet, Southeast China	late Aptian to Albian	<i>Shajia tibetica</i>	This paper
2	Ladakh, North India	late Aptian	<i>S. tibetica</i>  (= ' <i>Polyconites</i> sp.')	Masse and Fenerci-Masse, 2017
3	Shukur, Nubra-Shyok valley, Ladakh, North India	latest Aptian to early Albian	<i>Horiopleura</i>  <i>haydeni</i>	Upadhyay, 2001, 2014
4	Yasin, Gilgit region, Kohistan, Northwest Pakistan	latest Aptian to middle Albian	<i>H. haydeni</i>	Douvill�, 1926; Rossi Ronchetti, 1965; Pudsey et al., 1985; Sha and Cestari, 2016
5	Adi Gar Mt. (about 30 km N.N.W. from Spin Boldak) in Kandahar, Central Afghanistan	late Aptian	<i>H. haydeni</i>	Montenat et al., 1982
6	Khist hills, south of Khash Rud, Central Afghanistan	latest Aptian to middle Albian	<i>H. haydeni</i>	Rao et al., 2017






















## 1    **Highlights**

- 2    1. A new polyconitid rudist *Shajia tibetica* gen. et sp. nov., likely late Aptian to  
3        Albian in age, is described from the Langshan Formation of Nyima County, Tibet.
- 4    2. *Shajia* is externally comparable with *Horiopleura haydeni* Douvillé, 1926, but  
5        differs in its possession of an inwardly inclined, instead of outwardly facing,  
6        posterior myophore in the right valve
- 7    3. A single specimen from Ladakh that was assigned to *Polyconites?* sp. by Masse and  
8        Fenerci-Masse (2017), is revised and transferred to *Shajia*.
- 9    4. The *Shajia tibetica* /*Auroradiolites biconvexus* rudist association can be considered  
10        a regional variant of the Yasin fauna.

<sup>6</sup>Nanjing Center, China Geological Survey, Nanjing 210016, China; e-mail:

[pengbocc@163.com](mailto:pengbocc@163.com)<sup>†</sup>

<sup>7</sup>Sichuan Geological Survey, No. 25 North Renmin Road, Chengdu 610084, China; e-

mail: [986573254@qq.com](mailto:986573254@qq.com), [mzx0120@163.com](mailto:mzx0120@163.com)<sup>†</sup>